

# HIV-1 reverse transcriptase and antiviral drug resistance. Part 2

Kalyan Das and Eddy Arnold

Structures of RT and its complexes combined with biochemical and clinical data help in illuminating the molecular mechanisms of different drug-resistance mutations. The NRTI drugs that are used in combinations have different primary mutation sites. RT mutations that confer resistance to one drug can be hypersensitive to another RT drug. Structure of an RT-DNA-nevirapine complex revealed how NNRTI binding forbids RT from forming a polymerase competent complex. Collective knowledge about various mechanisms of drug resistance by RT has broader implications for understanding and targeting drug resistance in general. In Part 1, we discussed the role of RT in developing HIV-1 drug resistance, structural and functional states of RT, and the nucleoside/nucleotide analog (NRTI) and non-nucleoside (NNRTI) drugs used in treating HIV-1 infections. In this part, we discuss structural understanding of various mechanisms by which RT confers antiviral drug resistance.

## Addresses

Center for Advanced Biotechnology and Medicine (CABM), Department of Chemistry and Chemical Biology, Rutgers University, Piscataway, NJ 08854, USA

Corresponding author: Arnold, Eddy ([arnold@cabm.rutgers.edu](mailto:arnold@cabm.rutgers.edu))

**Current Opinion in Virology** 2013, **3**:119–128

This review comes from a themed issue on **Virus structure and function**

Edited by **Joseph Marcotrigiano** and **Felix Rey**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 19th April 2013

1879-6257/\$ – see front matter, © 2013 Elsevier B.V. All rights reserved.

<http://dx.doi.org/10.1016/j.coviro.2013.03.014>


## NRTI resistance by discrimination/exclusion

The dNTP substrates and analogs enter the open dNTP-binding cleft (**Figure 1a**), and the correct dNTP is selected for incorporation based on its complementarity (Watson–Crick base pairing) with the template overhang base (**Figure 1b,c**). The entire process starting from the entry of dNTP to the incorporation of a nucleotide is continuous and fast, although sorting of incorrectly paired bases limits the rate of incorporation. An NRTI-TP competes for RT binding with dNTPs; however, an NRTI-TP does not inhibit dNTP binding, and usually both bind RT at approximately micromolar affinity. HIV-1 inhibition by an NRTI occurs primarily when the NRTI is incorporated into the DNA primer. An NRTI-TP usually has modifications of its deoxyribose part when compared with a dNTP.

A dNTP (or NRTI-TP) undergoes association and dissociation of several molecular interactions with RT through the path starting from dNTP recognition by RT to nucleotide addition into a DNA primer strand. In the process, the distinct structural features of an NRTI-TP can be recognized by a specific mutation or sets of mutations that discriminate the drug from the analogous dNTP. For example, the active site mutation M184V/I causes high-level resistance to 3TC and FTC. A  $\beta$ -branched valine or isoleucine at the position 184 results in steric hindrance with the  $\beta$ -L-pseudo-ribose ring and thereby discriminates 3TC-TP (or FTC-TP) from dCTP [1,2]. Both 3TC and FTC inhibit hepatitis B, and the virus develops resistance to these drugs via analogous mutations in hepatitis B polymerase [3]. The HIV-1 RT mutation K65R emerges in response to treatments with TDF, ABC, ddI or d4T [4–6], and the mutation occurs at a higher frequency in subtype C HIV-1 [7]. The  $\beta$ 3- $\beta$ 4 residue K65 interacts with the  $\gamma$ -phosphate of a dNTP in RT-DNA-dNTP ternary complex structures. Structures of K65R RT-DNA-dATP (or TFV-DP) [8<sup>\*</sup>] revealed that the guanidinium side-chain functionality introduced by the K65R mutation stacks with the guanidinium group of R72, a key conserved residue that is involved in dNTP binding. The K65R-R72 guanidinium platform (**Figure 1d**) helps discriminate TFV-DP from dATP. Both wild-type and the mutant RT incorporated dATP and dATP- $\alpha$ -S at a similar rate indicating no elemental effect by K65R mutation, that is, the mutant RT discriminates at a conformational step rather than at the chemical step of NRTI incorporation [8<sup>\*</sup>].

The positioning of the sugar moiety of a dNTP/NRTI-TP at the N site is primarily defined by its interactions with amino acid residues R72, Y115, and Q151. The conserved residue R72 helps dNTP binding by stacking with the base and interacting with the  $\alpha$ -phosphate. The mutation Y115F emerged in response to ABC in mono- or combination therapies [9]. The mutation Y115F in combination with K65R decreases TDF susceptibility [10]. Residue Y115 is positioned underneath the sugar moiety and supports dNTP binding (**Figure 1d**). The Y115F mutation, which would retain its role in dNTP binding, acquires the ability of NRTI discrimination; wild-type hepatitis B polymerase has a phenylalanine at the equivalent position [3]. RT mutation at a nearby position, L74V, confers resistance to ddI [11,12]. Residue L74 is positioned under the base of the first template overhang and supports base-pair formation with an incoming dNTP (**Figure 1e**). Modeling of the L74V mutation [8<sup>\*</sup>] suggests that the mutation apparently weakens the template:dNTP base-pairing and/or causes repositioning of

Figure 1



NRTI-TP (or dNTP) binding and mechanisms of NRTI resistance. **(a)** Positioning of an AZTTP molecule in an open dNTP-binding cleft of RT (represented by the electrostatic potential surface) in an RT-DNA complex [68\*\*]. **(b)** RT-DNA-dTTP polymerase complex before catalysis was modeled using structural information from RT-DNA-dTTP complex [1], RT-ATP (a non-productive complex) [69], and RT-DNA-AZTTP complexes [68\*\*]; the surrounding residues (cyan) can mutate to confer resistance to NRTIs. **(c)** A schematic representation of dNTP binding with the help of base-pairing, base-stacking, interaction with RT, and metal chelation; the catalytic reaction of DNA polymerization proceeds with an octahedral coordination environment for two  $Mg^{2+}$  ions. **(d)** An active-site superposition of K65R RT-DNA-dATP [8\*] and EEM RT-DNA-AZTppppA complexes [23\*\*] shows the relative locations of three distinct sites that confer resistance to three classes of NRTIs (3TC/FTC, TDF, and AZT). The 3TC-resistance mutation introduces a  $\beta$ -branch, TDF-resistance mutation K65R forms the K65R-R72 guanidinium platform, and primary EEMs T215Y and K70R help binding of an ATP molecule as the excision agent; mutations at one site, in general, are incompatible with mutations at the other two sites which provides a partial structural/biochemical basis for synergy between NRTIs. **(e)** Molecular surface representing the support to template-dNTP base pair provided by R72 and L74 side-chains. **(f)** An L74V mutation would disrupt the platform, and the mutation is incompatible with the K65R induced K65R + R72 part of the platform. **(g)** A schematic representation of catalytic reaction of polymerization versus excision by HIV-1 RT [23\*\*]. **(h)** Binding of AZTppppA, the ATP-mediated excision product of AZTTP, to HIV-1 RT (shown as green molecular surface) in complex with DNA [23\*\*].

dNTP by reducing the support to the template base (Figure 1f).

Residue Q151 interacts with the guanidinium group of R72 [8\*]. The fact that the Q151 side chain in wild-type RT is unable to discriminate between a dNTP and a dideoxynucleoside-TP suggests that there may not be a

stable interaction between Q151 and the 3'-OH of a dNTP. In contrast, Q151M mutation can cause resistance to dideoxynucleoside drugs. Q151M and four mutations (A62V, V75I, F77L, and F116Y), known as the 'Q151M complex,' co-emerged as a set in response to AZT combined with dideoxynucleoside drugs [13,14]. The Q151M complex causes resistance to almost all NRTIs except

3TC/FTC and TDF, and the viral strains carrying the mutations are termed multidrug-resistant (MDR) viruses. The specific structural basis of NRTI resistance caused by Q151M or Q151M complex RT has not been characterized; however, it has been shown biochemically that Q151M mutation discriminates NRTI-TPs [15–17], and the accompanying four mutations in the Q151M complex enhance the discrimination ability and viral fitness [13,18]. RTs carrying Q151M mutation develop resistance by discriminating NRTI-TPs from dNTPs before incorporation [19].

### NRTI resistance by excision

A set of RT mutations (M41L, D67N, K70R, L210W, T215Y, and K219Q), known as thymidine analog mutations (TAMs) or AZT-resistance mutations (AZTr) [20], emerged in patients as distinctive mutations leading to AZT resistance [20]. Later, these mutations were biochemically shown to help excise AZT after the drug was incorporated into the DNA primer [21,22]. Excision is the primary mechanism by which RT develops resistance to AZT. Subsequently, several additional mutations were found to assist excision of AZT or other NRTIs, and this list of mutations can be grouped as Excision Enhancing Mutations (EEMs) [23<sup>••</sup>]. RT has the ability to excise either a normal nucleotide or an NRTI from the 3'-end of a DNA primer by pyrophosphorolysis, a reverse reaction of DNA polymerization (Figure 1g). Wild-type RT can excise the primer terminal nucleotide using a pyrophosphate or an ATP molecule as the excision agent; however, an RT carrying EEMs (TAMs) has a significantly enhanced excision ability using cellular ATP molecules as the excision agent [24,25]. The mutations facilitate ATP-dependent excision, generating AZTppppA (Figure 1g) as the reaction product. Crystal structures of RT-DNA-AZTppppA and related complexes illuminate the molecular mechanism of ATP-mediated excision of AZT (Figure 1g,h) [23<sup>••</sup>,26]. In agreement with an earlier hypothesis [25], the RT-DNA-AZTppppA structures revealed that the mutations help create an ATP-binding pocket adjacent to the dNTP-binding cleft. The T215Y mutated aromatic side chain stacks with the base while the K70R mutated side chain forms polar interactions with the  $\alpha$ -phosphate and 3'-OH of ATP. The structural information revealed that the primary AZT-resistance mutations K70R and T215Y [27,28] play the most significant roles in ATP binding. The  $\beta$ - $\gamma$ -diphosphate moiety of ATP mimics a pyrophosphate that chelates the two active site  $Mg^{2+}$  ions and triggers pyrophosphorolysis by RT to remove AZT-MP from the DNA primer terminus. The mutation-induced pocket has the potential to bind all NTPs and dNTPs; however, ATP is primarily used as the excision agent *in vivo* because: firstly, pyrophosphate is present at relatively low concentration in a cell; secondly, a dNTP would favor binding at the N site to form a polymerase catalytic complex rather than an excision complex;

thirdly, an NTP with a purine base would have more extensive stacking with T215Y than one with a pyrimidine base; GTP was also shown to be an efficient excision substrate [22]; and fourthly, the cellular concentration of ATP is about a order magnitude higher than that of GTP [29]. The remaining EEMs play supportive roles in enhancing excision and improving viral fitness on top of the primary EEM backgrounds [23<sup>••</sup>,30,31]. The mutations D67N, and K219Q [32–34] arise on top of the K70R background, whereas M41L and L210W mutations are associated with T215Y mutation [35,36]. The T69 insertions in the  $\beta$ 3– $\beta$ 4 fingers loop in combination with TAMs allow RT to excise a broader range of NRTIs [37–40]. Also, RT containing the  $\Delta$ 67 deletion enhances ATP-mediated excision of AZT on top of the TAM background [41,42]; however, the  $\Delta$ 67 RT does not excise a wider range of NRTIs.

In a fascinating parallel development, the hepatitis C (HCV) virus RNA-directed RNA polymerase (NS5B) has been shown to be capable of catalyzing NTP-mediated nucleotide excision [43,44<sup>••</sup>]. Analogous to the NTP-mediated nucleotide excision reaction by HIV-1 RT, that produces a dinucleoside tetraphosphate, the excision product by HCV NS5B (RNA polymerase) has ribonucleosides on both ends. Using a kinetic study, Jin et al. described that HCV polymerase excises chain-terminators and mismatched NMPs from 3'-end of a growing RNA primer [44<sup>••</sup>]. This result suggests that HCV and other polymerases may be using excision as a general fidelity mechanism. A lower rate of misincorporation by HIV-1 RT *in vivo* than free RT, as discussed in Part 1, may indicate that RT in infected cell conceivably may be using excision to enhance fidelity.

### Cooperative/antagonistic mutations and NRTI combinations in therapy

In general, members of a class of inhibitors such as NNRTIs or PIs, that share a common binding site and mode of inhibition, do not show synergistic inhibition if combined. In contrast, many combinations of NRTIs have synergistic impact in inhibiting HIV-1 in patients. The NRTI synergies were found empirically from virological investigations and from clinical data; however, effective modeling and analysis of empirical data [45<sup>••</sup>,46<sup>•</sup>], and biochemical and structural studies help in understanding the complex synergistic relationships and their molecular basis. From a simplified structural and biochemical perspective, a synergistic relationship between two NRTI drugs can be influenced by the antagonistic relationship among the primary resistance mutations to the NRTIs. Primary NRTI-resistance exclusion mutations appear along the dNTP-binding track, or EEMs emerge at sites adjacent to the dNTP-binding track to facilitate excision. The sites and molecular mechanisms of resistance to one subclass of NRTIs are different than those to another subclass of NRTIs [8<sup>•</sup>];



the primary resistance mutation sites are distinct for 3TC, TDF, and AZT (Figure 1d). Resistance mutations to one subclass of NRTI can be positively or negatively associated with the resistance mutations to another subclass. The K65R + M184V double mutant is primarily detected in patients treated with a TDF, 3TC, and ABC combination [47]. The double mutant has increased resistance to 3TC and FTC, decreased rate of nucleotide incorporation, and increased susceptibility to TDF [17,48]. The  $\beta$ -branch introduced by the M184V mutation, and the K65R-induced R72-K65R guanidinium platform, form two walls on either side of the sugar moiety of a dNTP/NRTI-TP at the N site (Figure 1d). Thereby, the K65R + M184V mutations create structural constraints [8<sup>•</sup>] that firstly, still can accommodate the compact acyclic methoxypropyl moiety of TFV-DP, secondly, add restraint to dNTP binding that would slow down nucleotide incorporation, and thirdly, are highly discriminatory against the  $\beta$ -L-pseudo-ribose ring of 3TC (or FTC)-TP. Even though both K65R and M184V are exclusion mutations that co-emerge, a lower replication fitness and increased susceptibility to TDF may account for the clinical benefit of combining TDF and 3TC (or FTC).

Among other exclusion mutations, K65R is negatively associated with L74V [49,50] or most EEMs [51], and the combinations emerge less frequently. The mutations K65R and L74V are incompatible because the K65R-R72 platform and L74V-modified surface do not provide optimal structural support to dNTP-template base pairing [8<sup>•</sup>] (Figure 1e,f); K65R + L74V double mutant RT has a significantly reduced rate of nucleotide incorporation [49]. Also, L74V mutation decreases AZT excision [11,52]; probably by perturbing the positioning of AZT (at the DNA primer end) at the N site even when EEMs facilitate ATP binding. The multidrug-resistant Q151M complex in association with other exclusion mutations K65R or K70T/S/Q increases resistance to TDF while remaining resistant to dideoxynucleosides [53,54].

In general, exclusion mutations and EEMs are antagonistic [55–57]. Structurally, the K65R + R72 platform would interfere with proper positioning of the  $\beta$ - $\gamma$ -pyrophosphate moiety of ATP, which is essential for excision [8<sup>•</sup>] (Figure 1d). Similarly, K70R and K65R would perturb the positioning of the triphosphate moiety of a dNTP and interfere with polymerization. Unlike the EEM K70R, all other mutations K70E/Q/S/T at position 70 function as exclusion mutations because these mutated side chains would not help ATP binding; however, the mutations may reposition or sense the  $\gamma$ -phosphate of an incoming NRTI-TP differently from that of a dNTP and invoke NRTI resistance by an exclusion mechanism. The mutation K70E is also antagonistic with EEMs [58], because the negatively charged shorter side chain of K70E is likely to interfere with the binding or positioning


of ATP (Figure 1d,h). M184V/I mutation is antagonistic with excision. M184V/I mutation, which reduces the binding of 3TC (or FTC) at the N site, also may hinder the positioning of an NRTI-terminated 3' end of a DNA primer at the N site for excision, that is, 3TC resistance enhances AZT susceptibility which favors the synergistic impact of combining AZT and 3TC [59].

## RT inhibition by NNRTIs

RT inhibition by NNRTIs is indirect. Early structures of RT and RT-NNRTI complexes showed that NNRTI binding traps RT in a rigid conformational state with an open nucleic acid-binding cleft; however, the impact of NNRTI binding on the binding of nucleic acid or dNTP, and above all, the mechanism of NNRTI inhibition of DNA polymerization remained elusive. Pre-steady-state and steady-state kinetic experiments in the mid-1990s suggested that the binding of an NNRTI inhibits the chemical step of DNA polymerization while dsDNA and dNTP remain bound to RT [60,61]. A single-molecule FRET study revealed that RT frequently flips and slides over a double-stranded nucleic acid substrate [62<sup>••</sup>,63]; binding of dNTP stabilized the RT-DNA complex in a polymerase-competent mode while binding of an NNRTI induced destabilizing effects causing increased dissociation/association of RT with a double-stranded nucleic acid. The postulated mechanisms of inhibition by an NNRTI were: firstly, restriction of thumb mobility [64], secondly, distortion of the catalytic triad [65] that would block the chemical step of DNA polymerization by RT [60], thirdly, blockage of conformational state transition and not the chemical step leading to nucleotide incorporation [66], fourthly, repositioning of the primer grip [67], and fifthly, loosening of the thumb and fingers clamp [62<sup>••</sup>]. Multiple mechanistic hypotheses evolved because NNRTI binding causes multiple structural and conformational changes in RT, and the direct and indirect contributions of individual changes toward the NNRTI inhibition were not clear.

The hinge motion between the thumb and palm is essential for the translocation of RT along the nucleic acid following each nucleotide incorporation. The breathing space that is transiently created traps an NNRTI. Upon binding of an NNRTI, RT loses its conformational mobility to carry out nucleotide addition and translocation. The NNRTI pocket does not exist in any RT structure that does not contain a bound NNRTI. A direct consequence of NNRTI binding is the opening of the NNRTI pocket to accommodate the inhibitor. The pocket formation requires the  $\beta$ 12– $\beta$ 13– $\beta$ 14 sheet, which contains the 'primer grip', to move away from the  $\beta$ 6– $\beta$ 10– $\beta$ 9 sheet which contains the catalytic triad (D110, D185, and D186). The recent structure of RT-DNA-nevirapine complex provides a snapshot of the effects of an NNRTI on DNA polymerization [68<sup>••</sup>] (Figure 2a,b). The binding of nevirapine shifts the primer grip which

Figure 2



Structural basis for the inhibition of DNA polymerization by an NNRTI. (a) Structure of RT-DNA-AZTTP ternary complex obtained by soaking AZTTP into crystals of RT-DNA complex. (b) Soaking of nevirapine into the crystal created the NNRTI pocket, repositioning the 'primer grip' (on the  $\beta 12$ - $\beta 13$ - $\beta 14$  sheet) that moved the primer terminus away from the polymerase active site. (c) Electrostatic potential surface of RT bound to DNA and nevirapine. The crystallization experiments and structures showed an open dNTP-binding cleft into which dNTPs/NNRTI-TPs can enter; however, structural perturbation by NNRTI binding did not allow a dNTP to chelate metals and form base-pairing and base-stacking interactions. (d) These structural constraints preclude the formation of an RT-DNA-dNTP polymerase competent (P) complex, rather forms a non-productive (P') complex in the presence of an NNRTI — a structural basis for NNRTI inhibition [68\*\*].

concomitantly displaces the primer 3'-end by  $\sim 5.5$  Å away from its position at the polymerase active site. The interaction between the template-primer and the polymerase domain of RT is decreased upon nevirapine binding, which correlates with the earlier observation by a single-molecule study [62\*\*]. The fingers subdomain has an open conformation that would allow the entry of dNTPs (Figure 2c); however, the repositioned template-primer would not permit the base-pairing or base-stacking to support the binding of a dNTP (Figure 1c) at

the N site. This appears to be a primary reason why ordered binding of dNTP (or analog) to an RT-DNA complex was not observed in the presence of nevirapine whereas RT-DNA-dNTP (or analog) complexes could be formed when no NNRTI was present [68\*\*]. These experiments were carried out in crystals that permitted rearrangements of the polymerase domain upon binding of a dNTP/analog or an NNRTI. Our attempts to form an RT-DNA-nevirapine-AZTTP complex yielded the structure of only RT-DNA-nevirapine complex, and

no well-defined binding of AZTTP at the polymerase active site when nevirapine is bound to RT [68<sup>••</sup>]. The structural information and published biochemical/biophysical results suggest that dNTPs may enter the dNTP-binding cleft (Figure 2c), interact with RT, and may induce conformational changes of RT like closing of the fingers; however, formation of a catalytically relevant RT-DNA-dNTP complex [1] would not be permitted when an NNRTI is bound. The repositioning of the template-primer by nevirapine binding disfavors the base-pairing or base-stacking of a dNTP at the N site. In addition to the base-pairing and base-stacking, metal chelation at the active site also contributes toward the binding of a dNTP substrate in a catalytically competent RT-DNA-dNTP complex (Figure 1b,c). None of the structures of RT-NNRTI binary complexes or the RT-DNA-nevirapine complex had any metal ion present at the polymerase active site, suggesting that a potential distortion of the catalytic site [65,69] by an NNRTI also forbids ordered binding of the triphosphate moiety of a dNTP because of the loss of Mg<sup>2+</sup> ion chelation at the active site. Thereby, NNRTI binding prevents RT from achieving a conformational state of RT-DNA (or RT-DNA/RNA) complexed with dNTP that would be required for catalysis [66]. Additionally, the structure of RT-DNA-nevirapine complex showed that the thumb restriction [64] is induced by the primer grip repositioning, and the thumb and fingers clamp is loosened [62<sup>••</sup>] by the reduced interactions between the polymerase domain of RT and DNA. All of these effects of NNRTI binding force RT into a structurally and catalytically non-competent complex for dNTP binding at the N site and for polymerization (Figure 2d).

### NNRTI resistance

Almost all of the NNRTI-binding pocket residues can mutate, and most of the mutations can confer resistance to NNRTIs. Some of the observed NNRTI-resistance mutations are L100I, K101E, V106A, K103N, V179D, Y181C, Y188L, G190A, and E138K (p51); K103N and Y181C are the most commonly observed mutations in patients exposed to NNRTI treatments. Residues K101, K103, and E138 (p51) are located at the rim of the pocket entrance for most NNRTIs. The remaining residues directly interact with an NNRTI in the pocket. The pocket mutations can cause firstly, loss of aromatic ring stacking interactions (Y181C or Y188L), secondly, steric hindrance (L100I or G190A/S), and thirdly, alteration of hydrophobic interactions (V106A or V179D). Impacts of NNRTI-pocket alteration are relatively severe on the rigid first-generation NNRTIs, for example, high level of resistance by Y181C to nevirapine. Substitution of G190 fills the central part of the pocket by which mutations at position 190 would reduce the binding of most NNRTIs; however, larger substitutions like G190E/Q severely impair the replication capacity of RT [70,71]; G190A is the most frequently observed mutation at position 190

because G190A mutation has the least impact on RT polymerase activity. A large side chain at position 190 would partly occupy the NNRTI-binding pocket and could conceivably function as a pseudo-NNRTI in reducing RT activity. A fitness cost is associated with certain pocket mutations or combinations of mutations if the virus has to develop resistance to more effective NNRTIs. Such NNRTIs will have to remain potent against all common NNRTI-resistance mutations that have minimum adverse impact on the viral replication.

Rim mutations at K101, K103, or E138 (p51) generally do not alter the bound states of NNRTIs. Two recent clinical studies showed that K103N and E138K mutations are predominantly associated with treatment failure of the efavirenz and rilpivirine, respectively, when combined with TDF and FTC [72<sup>•</sup>,73<sup>•</sup>]; M184V was the NRTI-resistance mutation that co-emerged in both arms of the study. The study indicated a comparable response by 25 mg/day of rilpivirine or 600 mg/day of efavirenz; rilpivirine was better tolerated, yet had a higher rate of virological failure. Efavirenz is the most widely used NNRTI, and K103N is the characteristic clinical resistance mutation that emerged upon treatments with efavirenz. In fact, the K103N mutation confers an almost uniform level of cross-resistance to most NNRTIs. In crystal structures of wild-type RT, the K103 side chain points outward and does not interact with a bound NNRTI. The K103N mutation apparently restricts access to the pocket for most NNRTIs. In absence of an NNRTI, the shrunken pocket accommodated a water molecule [74], whereas the K103N/Y181C mutant RT apo structure revealed that the solvent molecule was replaced by a sodium ion that had a distorted octahedral coordination involving the side chains of N103 and Y188, and the main-chain carbonyl oxygen of K101 and amide group of N103 [69]; the side chains of N103 and Y188 were also in position to make a hydrogen-bonding interaction. The difference in the energetic cost required for pocket formation by displacing the secluded water versus ion, or the extra hydrogen bond, may contribute to NNRTI resistance; however, the exact mechanism by which the K103N mutation develops resistance remains unclear. In rilpivirine, which remains effective against K103N mutation, the central pyrimidine ring interacts with the side chain of N103 [75]. A computational study using Monte Carlo/free energy perturbation calculations had suggested that NNRTIs that would form water-mediated hydrogen bonds at the entrance to the pocket, in particular with E138 (p51), would retain potency against K103N mutation [76].

The rim mutations K101E or E138K (p51) appear to have effects on NNRTI binding analogous to that of K103N. In wild-type RT, the side chains of E138 (p51) and K101 form a hydrogen bond that is lost in E138K or K101E mutant RTs [77]; the loss of the hydrogen bond was assumed to be the primary cause of NNRTI resistance



by E138K or K101E. If so, a mutant RT containing both E138K and K101E would restore the hydrogen bond between the two residues, and the double mutant RT would restore efficacy of NNRTIs. However, the double mutant showed a high level of NNRTI resistance and the E138D mutant RT that would not form the hydrogen bond with K101 remained NNRTI-susceptible just by retaining the charge of E/D138 side chain [78], which downweighs the suggested critical role of the hydrogen bond between K101 and E138 on NNRTI binding. A recent kinetic study indicated that rilpivirine had a higher rate of dissociation from E138K (p51) mutant versus wild-type RT [79<sup>\*</sup>]. The cumulative biochemical and structural knowledge so far indicates that the rim mutations K103N and E138K primarily alter the association and dissociation of NNRTIs with RT. Even though E138K itself only causes a low level of resistance to rilpivirine, addition of M184V/I mutation enhances rilpivirine resistance, and E138K enhances the fitness of the virus on the M184V/I background, which reveals a synergistic relationship between the NNRTI- and NRTI-resistance mutations that benefits the virus [72<sup>\*</sup>]. In contrary, the NNRTI-resistance mutation Y181C on a background of NRTI-resistance mutations decreases ATP binding as the excision substrate [80], indicating an antagonistic relationship.

### Use of NRTIs and NNRTIs in combinations

Effective combinations of HIV-1 drugs are being optimized empirically based on clinical data. One or more NRTIs are used in almost all current clinical drug cocktails, and many combinations of NRTIs are synergistic. RT mutations that emerge along the dNTP-binding cleft to discriminate NRTI-TPs or to enhance excision of an NRTI are often incompatible for RT fitness, or mutations that develop resistance to one NRTI enhance sensitivity to another – among the molecular bases for the long-term clinical utility of NRTIs in combinations. Because of the different inhibition mechanisms, combinations of NNRTIs and NRTIs are effective in reducing viral replication. NNRTIs have the advantage of high specificity and consequently low side effects. An NNRTI is primarily used in combination with NRTI(s) in first-line therapies against HIV-1 infections; however, higher level of treatment failure has been observed for patients carrying traces of NNRTI-resistance mutations [81<sup>\*</sup>]. NNRTIs are also important for blocking mother-to-child transmission, and NNRTIs are currently under clinical investigations as potential microbicides to prevent sexual transmission of HIV-1 [82]. The relationships among RT mutations are complex and are still evolving [46<sup>\*</sup>]. New resistance patterns [83,84] emerge with the use of new drugs and combinations. No single experimental approach or analysis fully explains the complex phenomena associated with HIV-1 infection and drug resistance. Multiple experimental platforms, analysis of experimental data across the platforms, and their relationships with clinical observations are continually helping to unfold

new details on the virus and antiviral resistance. Frequent evolution of new resistance mutation patterns and emerging side effects from long-term use of drugs underscore the constant need for new HIV-1 drugs.

### Acknowledgements

EA is grateful to the National Institutes of Health for support from grants R37 AI 27690 (MERIT Award) and P50 GM103368.

### References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
  - of outstanding interest
- Huang H, Chopra R, Verdine GL, Harrison SC: **Structure of a covalently trapped catalytic complex of HIV-1 reverse transcriptase: implications for drug resistance.** *Science* 1998, **282**:1669-1675.
  - Sarafianos SG, Das K, Clark AD Jr, Ding J, Boyer PL, Hughes SH, Arnold E: **Lamivudine (3TC) resistance in HIV-1 reverse transcriptase involves steric hindrance with  $\beta$ -branched amino acids.** *Proc Natl Acad Sci U S A* 1999, **96**:10027-10032.
  - Das K, Xiong X, Yang H, Westland CE, Gibbs CS, Sarafianos SG, Arnold E: **Molecular modeling and biochemical characterization reveal the mechanism of hepatitis B virus polymerase resistance to lamivudine (3TC) and emtricitabine (FTC).** *J Virol* 2001, **75**:4771-4779.
  - Winters MA, Shafer RW, Jellinger RA, Mamtara G, Gingeras T, Merigan TC: **Human immunodeficiency virus type 1 reverse transcriptase genotype and drug susceptibility changes in infected individuals receiving dideoxyinosine monotherapy for 1 to 2 years.** *Antimicrob Agents Chemother* 1997, **41**:757-762.
  - Harrigan PR, Stone C, Griffin P, Najera I, Bloor S, Kemp S, Tisdale M, Larder B: **Resistance profile of the human immunodeficiency virus type 1 reverse transcriptase inhibitor abacavir (1592U89) after monotherapy and combination therapy.** CNA2001 Investigative Group. *J Infect Dis* 2000, **181**:912-920.
  - Margot NA, Lu B, Cheng A, Miller MD: **Resistance development over 144 weeks in treatment-naïve patients receiving tenofovir disoproxil fumarate or stavudine with lamivudine and efavirenz in Study 903.** *HIV Med* 2006, **7**:442-450.
  - Coutsinos D, Invernizzi CF, Xu H, Moisi D, Oliveira M, Brenner BG, Wainberg MA: **Template usage is responsible for the preferential acquisition of the K65R reverse transcriptase mutation in subtype C variants of human immunodeficiency virus type 1.** *J Virol* 2009, **83**:2029-2033.
  - Das K, Bandwar RP, White KL, Feng JY, Sarafianos SG, Tuske S, Tu X, Clark AD Jr, Boyer PL, Hou X *et al.*: **Structural basis for the role of the K65R mutation in HIV-1 reverse transcriptase polymerization, excision antagonism, and tenofovir resistance.** *J Biol Chem* 2009, **284**:35092-35100.
  - Structures of K65R RT/DNA/dATP (or TFV-DP) showed how K65R mutation forms a R72 + K65R platform that discriminates the drug from dATP. The distinct locations of 3TC-, TDF-, and AZT-resistance mutation sites on RT provide a structural basis for synergistic benefits of NRTI drug combinations in suppressing resistance mutations.
  - Miller V, Ait-Khaled M, Stone C, Griffin P, Mesogiti D, Cutrell A, Harrigan R, Staszewski S, Katlama C, Pearce G *et al.*: **HIV-1 reverse transcriptase (RT) genotype and susceptibility to RT inhibitors during abacavir monotherapy and combination therapy.** *AIDS* 2000, **14**:163-171.
  - Stone C, Ait-Khaled M, Craig C, Griffin P, Tisdale M: **Human immunodeficiency virus type 1 reverse transcriptase mutation selection during in vitro exposure to tenofovir alone or combined with abacavir or lamivudine.** *Antimicrob Agents Chemother* 2004, **48**:1413-1415.

11. St Clair MH, Martin JL, Tudor-Williams G, Bach MC, Vavro CL, King DM, Kellam P, Kemp SD, Larder BA: **Resistance to ddl and sensitivity to AZT induced by a mutation in HIV-1 reverse transcriptase.** *Science* 1991, **253**:1557-1559.
12. Martin JL, Wilson JE, Haynes RL, Furman PA: **Mechanism of resistance of human immunodeficiency virus type 1 to 2',3'-dideoxyinosine.** *Proc Natl Acad Sci U S A* 1993, **90**:6135-6139.
13. Shirasaka T, Kavlick MF, Ueno T, Gao WY, Kojima E, Alcaide ML, Choekijchai S, Roy BM, Arnold E, Yarchoan R *et al.*: **Emergence of human immunodeficiency virus type 1 variants with resistance to multiple dideoxynucleosides in patients receiving therapy with dideoxynucleosides.** *Proc Natl Acad Sci U S A* 1995, **92**:2398-2402.
14. Kavlick MF, Wyvill K, Yarchoan R, Mitsuya H: **Emergence of multi-dideoxynucleoside-resistant human immunodeficiency virus type 1 variants, viral sequence variation, and disease progression in patients receiving antiretroviral chemotherapy.** *J Infect Dis* 1998, **177**:1506-1513.
15. Kaushik N, Talele TT, Pandey PK, Harris D, Yadav PN, Pandey VN: **Role of glutamine 151 of human immunodeficiency virus type-1 reverse transcriptase in substrate selection as assessed by site-directed mutagenesis.** *Biochemistry* 2000, **39**:2912-2920.
16. Deval J, Selmi B, Boretto J, Eglhoff MP, Guerreiro C, Sarfati S, Canard B: **The molecular mechanism of multidrug resistance by the Q151M human immunodeficiency virus type 1 reverse transcriptase and its suppression using alpha-boranophosphate nucleotide analogues.** *J Biol Chem* 2002, **277**:42097-42104.
17. Deval J, White KL, Miller MD, Parkin NT, Courcambecq J, Halfon P, Selmi B, Boretto J, Canard B: **Mechanistic basis for reduced viral and enzymatic fitness of HIV-1 reverse transcriptase containing both K65R and M184V mutations.** *J Biol Chem* 2004, **279**:509-516.
18. Kosalaraksa P, Kavlick MF, Maroun V, Le R, Mitsuya H: **Comparative fitness of multi-dideoxynucleoside-resistant human immunodeficiency virus type 1 (HIV-1) in an in vitro competitive HIV-1 replication assay.** *J Virol* 1999, **73**:5356-5363.
19. Ray AS, Basavapathruni A, Anderson KS: **Mechanistic studies to understand the progressive development of resistance in human immunodeficiency virus type 1 reverse transcriptase to abacavir.** *J Biol Chem* 2002, **277**:40479-40490.
20. Larder BA, Kemp SD: **Multiple mutations in HIV-1 reverse transcriptase confer high-level resistance to zidovudine (AZT).** *Science* 1989, **246**:1155-1158.
21. Arion D, Kaushik N, McCormick S, Borkow G, Parniak MA: **Phenotypic mechanism of HIV-1 resistance to 3'-azido-3'-deoxythymidine (AZT): increased polymerization processivity and enhanced sensitivity to pyrophosphate of the mutant viral reverse transcriptase.** *Biochemistry* 1998, **37**:15908-15917.
22. Meyer PR, Matsuura SE, So AG, Scott WA: **Unblocking of chain-terminated primer by HIV-1 reverse transcriptase through a nucleotide-dependent mechanism.** *Proc Natl Acad Sci U S A* 1998, **95**:13471-13476.
23. Tu X, Das K, Han Q, Bauman JD, Clark AD Jr, Hou X, Frenkel YV, Gaffney BL, Jones RA, Boyer PL *et al.*: **Structural basis of HIV-1 resistance to AZT by excision.** *Nat Struct Mol Biol* 2010, **17**:1202-1209.
- Structure of TAM RT/DNA/AZTppppA (the ATP-mediated excision product of AZT) and related complexes revealed how TAM mutations help create a pocket for ATP to bind and excise NRTI (or nucleotide). Roles of the T215Y/F and K70R mutations in binding the ATP were visualized.
24. Meyer PR, Matsuura SE, Mian AM, So AG, Scott WA: **A mechanism of AZT resistance: an increase in nucleotide-dependent primer unblocking by mutant HIV-1 reverse transcriptase.** *Mol Cell* 1999, **4**:35-43.
25. Boyer PL, Sarafianos SG, Arnold E, Hughes SH: **Selective excision of AZTMP by drug-resistant human immunodeficiency virus reverse transcriptase.** *J Virol* 2001, **75**:4832-4842.
26. Scott WA: **Structures of reverse transcriptase pre- and post-excision complexes shed new light on HIV-1 AZT resistance.** *Viruses* 2011, **3**:20-25.
27. Boucher CA, O'Sullivan E, Mulder JW, Ramautarsing C, Kellam P, Darby G, Lange JM, Goudsmit J, Larder BA: **Ordered appearance of zidovudine resistance mutations during treatment of 18 human immunodeficiency virus-positive subjects.** *J Infect Dis* 1992, **165**:105-110.
28. Jeeninga RE, Keulen W, Boucher C, Sanders RW, Berkhout B: **Evolution of AZT resistance in HIV-1: the 41-70 intermediate that is not observed in vivo has a replication defect.** *Virology* 2001, **283**:294-305.
29. Jiang LJ, Maret W, Vallee BL: **The ATP-metallothionein complex.** *Proc Natl Acad Sci U S A* 1998, **95**:9146-9149.
30. Harrigan PR, Bloor S, Larder BA: **Relative replicative fitness of zidovudine-resistant human immunodeficiency virus type 1 isolates in vitro.** *J Virol* 1998, **72**:3773-3778.
31. Hu Z, Giguel F, Hatano H, Reid P, Lu J, Kuritzkes DR: **Fitness comparison of thymidine analog resistance pathways in human immunodeficiency virus type 1.** *J Virol* 2006, **80**:7020-7027.
32. Arion D, Parniak MA: **HIV resistance to zidovudine: the role of pyrophosphorolysis.** *Drug Resist Updat* 1999, **2**:91-95.
33. Yahi N, Tamalet C, Tourres C, Tivoli N, Ariasi F, Volot F, Gastaut JA, Gallais H, Moreau J, Fantini J: **Mutation patterns of the reverse transcriptase and protease genes in human immunodeficiency virus type 1-infected patients undergoing combination therapy: survey of 787 sequences.** *J Clin Microbiol* 1999, **37**:4099-4106.
34. Garcia-Lerma JG, MacInnes H, Bennett D, Weinstock H, Heneine W: **Transmitted human immunodeficiency virus type 1 carrying the D67N or K219Q/E mutation evolves rapidly to zidovudine resistance in vitro and shows a high replicative fitness in the presence of zidovudine.** *J Virol* 2004, **78**:7545-7552.
35. Harrigan PR, Kinghorn I, Bloor S, Kemp SD, Najera I, Kohli A, Larder BA: **Significance of amino acid variation at human immunodeficiency virus type 1 reverse transcriptase residue 210 for zidovudine susceptibility.** *J Virol* 1996, **70**:5930-5934.
36. Hooker DJ, Tachedjian G, Solomon AE, Gurusinge AD, Land S, Birch C, Anderson JL, Roy BM, Arnold E, Deacon NJ: **An in vivo mutation from leucine to tryptophan at position 210 in human immunodeficiency virus type 1 reverse transcriptase contributes to high-level resistance to 3'-azido-3'-deoxythymidine.** *J Virol* 1996, **70**:8010-8018.
37. Mas A, Parera M, Briones C, Soriano V, Martinez MA, Domingo E, Menendez-Arias L: **Role of a dipeptide insertion between codons 69 and 70 of HIV-1 reverse transcriptase in the mechanism of AZT resistance.** *EMBO J* 2000, **19**:5752-5761.
38. Boyer PL, Sarafianos SG, Arnold E, Hughes SH: **Nucleoside analog resistance caused by insertions in the fingers of human immunodeficiency virus type 1 reverse transcriptase involves ATP-mediated excision.** *J Virol* 2002, **76**:9143-9151.
39. Meyer PR, Lennerstrand J, Matsuura SE, Larder BA, Scott WA: **Effects of dipeptide insertions between codons 69 and 70 of human immunodeficiency virus type 1 reverse transcriptase on primer unblocking, deoxynucleoside triphosphate inhibition, and DNA chain elongation.** *J Virol* 2003, **77**:3871-3877.
40. White KL, Chen JM, Margot NA, Wrin T, Petropoulos CJ, Naeger LK, Swaminathan S, Miller MD: **Molecular mechanisms of tenofovir resistance conferred by human immunodeficiency virus type 1 reverse transcriptase containing a diserine insertion after residue 69 and multiple thymidine analog-associated mutations.** *Antimicrob Agents Chemother* 2004, **48**:992-1003.
41. Imamichi T, Berg SC, Imamichi H, Lopez JC, Metcalf JA, Falloon J, Lane HC: **Relative replication fitness of a high-level 3'-azido-3'-deoxythymidine-resistant variant of human immunodeficiency virus type 1 possessing an amino acid deletion at codon 67 and a novel substitution (Thr→Gly) at codon 69.** *J Virol* 2000, **74**:10958-10964.
42. Boyer PL, Imamichi T, Sarafianos SG, Arnold E, Hughes SH: **Effects of the Delta67 complex of mutations in human immunodeficiency virus type 1 reverse transcriptase on nucleoside analog excision.** *J Virol* 2004, **78**:9987-9997.



43. Deval J, Powdrill MH, D'Abramo CM, Cellai L, Gotte M: **Pyrophosphorolytic excision of nobligate chain terminators by hepatitis C virus NS5B polymerase.** *Antimicrob Agents Chemother* 2007, **51**:2920-2928.
  44. Jin Z, Leveque V, Ma H, Johnson KA, Klumpp K: **NTP-mediated nucleotide excision activity of hepatitis C virus RNA-dependent RNA polymerase.** *Proc Natl Acad Sci U S A* 2013, **110**:E348-E357.
- Using pre-steady-state kinetics, the study revealed that hepatitis C (HCV) RNA polymerase uses NTP-mediated excision to remove misincorporated nucleotides or chain terminators from the growing RNA strand. Although nucleotide excision by HCV polymerase is ~100-fold more efficient than that by HIV-1 RT, no excision-enhancing mutations have been reported yet in HCV polymerase.
45. Jilek BL, Zarr M, Sampah ME, Rabi SA, Bullen CK, Lai J, Shen L, Siliciano RF: **A quantitative basis for antiretroviral therapy for HIV-1 infection.** *Nat Med* 2012, **18**:446-451.
- The study analyzed dose-response curves for HIV-1 drugs from single-round infectivity assays in CD4+ T-cells. The analysis help rationalize several complex relationships among therapeutic combinations of drugs, and the approach developed may be critical in designing optimal drug combinations and in preclinical development of new drugs.
46. Melikian GL, Rhee SY, Taylor J, Fessel WJ, Kaufman D, Townner W, Troia-Cancio PV, Zolopa A, Robbins GK, Kagan R *et al.*: **Standardized comparison of the relative impacts of HIV-1 reverse transcriptase (RT) mutations on nucleoside RT inhibitor susceptibility.** *Antimicrob Agents Chemother* 2012, **56**:2305-2313.
- Using the PhenoSense assay, the study analyzed RT mutations in 1,273 HIV-1 isolates from patients with known treatment history. The analysis provides relative phenotypic effects of common NRTI-resistance mutations on each of the NRTIs.
47. Delaunay C, Brun-Vezinet F, Landman R, Collin G, Peytavin G, Trylesinski A, Flandre P, Miller M, Descamps D: **Comparative selection of the K65R and M184V/I mutations in human immunodeficiency virus type 1-infected patients enrolled in a trial of first-line triple-nucleoside analog therapy (Tonus IMEA 021).** *J Virol* 2005, **79**:9572-9578.
  48. White KL, Margot NA, Wrin T, Petropoulos CJ, Miller MD, Naeger LK: **Molecular mechanisms of resistance to human immunodeficiency virus type 1 with reverse transcriptase mutations K65R and K65R + M184V and their effects on enzyme function and viral replication capacity.** *Antimicrob Agents Chemother* 2002, **46**:3437-3446.
  49. Deval J, Navarro JM, Selmi B, Courcambeck J, Boretto J, Halfon P, Garrido-Urbani S, Sire J, Canard B: **A loss of viral replicative capacity correlates with altered DNA polymerization kinetics by the human immunodeficiency virus reverse transcriptase bearing the K65R and L74V dideoxynucleoside resistance substitutions.** *J Biol Chem* 2004, **279**:25489-25496.
  50. Sharma PL, Nurpeisov V, Lee K, Skaggs S, Di San Filippo CA, Schinazi RF: **Replication-dependent 65R→K mutation in human immunodeficiency virus type 1 reverse transcriptase double mutant K65R + L74V.** *Virology* 2004, **321**:222-234.
  51. McColl DJ, Chappey C, Parkin NT, Miller MD: **Prevalence, genotypic associations and phenotypic characterization of K65R, L74V and other HIV-1 RT resistance mutations in a commercial database.** *Antivir Ther* 2008, **13**:189-197.
  52. Miranda LR, Gotte M, Liang F, Kuritzkes DR: **The L74V mutation in human immunodeficiency virus type 1 reverse transcriptase counteracts enhanced excision of zidovudine monophosphate associated with thymidine analog resistance mutations.** *Antimicrob Agents Chemother* 2005, **49**:2648-2656.
  53. Van Laethem K, Pannecouque C, Vandamme AM: **Mutations at 65 and 70 within the context of a Q151M cluster in human immunodeficiency virus type 1 reverse transcriptase impact the susceptibility to the different nucleoside reverse transcriptase inhibitors in distinct ways.** *Infect Genet Evol* 2007, **7**:600-603.
  54. Hachiya A, Kodama EN, Schuckmann MM, Kirby KA, Michailidis E, Sakagami Y, Oka S, Singh K, Sarafianos SG: **K70Q adds high-level tenofovir resistance to 'Q151M complex' HIV reverse transcriptase through the enhanced discrimination mechanism.** *PLoS ONE* 2011, **6**:e16242.
  55. Wolf K, Walter H, Beerenwinkel N, Keulen W, Kaiser R, Hoffmann D, Lengauer T, Selbig J, Vandamme AM, Korn K *et al.*: **Tenofovir resistance and resensitization.** *Antimicrob Agents Chemother* 2003, **47**:3478-3484.
  56. White KL, Margot NA, Ly JK, Chen JM, Ray AS, Pavelko M, Wang R, McDermott M, Swaminathan S, Miller MD: **A combination of decreased NRTI incorporation and decreased excision determines the resistance profile of HIV-1 K65R RT.** *AIDS* 2005, **19**:1751-1760.
  57. Parikh UM, Bacheler L, Koontz D, Mellors JW: **The K65R mutation in human immunodeficiency virus type 1 reverse transcriptase exhibits bidirectional phenotypic antagonism with thymidine analog mutations.** *J Virol* 2006, **80**:4971-4977.
  58. Sluis-Cremer N, Sheen CW, Zelina S, Torres PS, Parikh UM, Mellors JW: **Molecular mechanism by which the K70E mutation in human immunodeficiency virus type 1 reverse transcriptase confers resistance to nucleoside reverse transcriptase inhibitors.** *Antimicrob Agents Chemother* 2007, **51**:48-53.
  59. Larder BA, Kemp SD, Harrigan PR: **Potential mechanism for sustained antiretroviral efficacy of AZT-3TC combination therapy.** *Science* 1995, **269**:696-699.
  60. Spence RA, Kati WM, Anderson KS, Johnson KA: **Mechanism of inhibition of HIV-1 reverse transcriptase by nonnucleoside inhibitors.** *Science* 1995, **267**:988-993.
  61. Rittinger K, Divita G, Goody RS: **Human immunodeficiency virus reverse transcriptase substrate-induced conformational changes and the mechanism of inhibition by nonnucleoside inhibitors.** *Proc Natl Acad Sci U S A* 1995, **92**:8046-8049.
  62. Abbondanzieri EA, Bokinsky G, Rausch JW, Zhang JX, Le Grice SF, Zhuang X: **Dynamic binding orientations direct activity of HIV reverse transcriptase.** *Nature* 2008, **453**:184-189.
- The study used a single-molecule FRET assay to investigate dynamic states of HIV-1 RT in complex with template-primer; RT was found to dynamically switch between two opposite modes of binding to double-stranded nucleic acid. Presence of dNTP or NNRTI impacts the relative population of the two distinct complexes.
63. Liu S, Abbondanzieri EA, Rausch JW, Le Grice SF, Zhuang X: **Slide into action: dynamic shuttling of HIV reverse transcriptase on nucleic acid substrates.** *Science* 2008, **322**:1092-1097.
  64. Kohlstaedt LA, Wang J, Friedman JM, Rice PA, Steitz TA: **Crystal structure at 3.5 Å resolution of HIV-1 reverse transcriptase complexed with an inhibitor.** *Science* 1992, **256**:1783-1790.
  65. Ren J, Esnouf R, Garman E, Somers D, Ross C, Kirby I, Keeling J, Darby G, Jones Y, Stuart D *et al.*: **High resolution structures of HIV-1 RT from four RT-inhibitor complexes.** *Nat Struct Biol* 1995, **2**:293-302.
  66. Xia Q, Radzio J, Anderson KS, Sluis-Cremer N: **Probing nonnucleoside inhibitor-induced active-site distortion in HIV-1 reverse transcriptase by transient kinetic analyses.** *Protein Sci* 2007, **16**:1728-1737.
  67. Das K, Ding J, Hsiou Y, Clark AD Jr, Moereels H, Koymans L, Andries K, Pauwels R, Janssen PA, Boyer PL *et al.*: **Crystal structures of 8-Cl and 9-Cl TIBO complexed with wild-type HIV-1 RT and 8-Cl TIBO complexed with the Tyr181Cys HIV-1 RT drug-resistant mutant.** *J Mol Biol* 1996, **264**:1085-1100.
  68. Das K, Martinez SE, Bauman JD, Arnold E: **HIV-1 reverse transcriptase complex with DNA and nevirapine reveals non-nucleoside inhibition mechanism.** *Nat Struct Mol Biol* 2012, **19**:253-259.
- The study showed that the RT-DNA complex in a crystal could bind an NNRTI or dNTP (NRTI-TP), but not both. The structure of RT-DNA-nevirapine complex illustrated that an NNRTI inhibits DNA polymerization by imposing conformational restrictions on the RT-DNA complex. The primer terminus is shifted in conjunction with the primer grip by more than 5 Å away from its position in catalytically competent conformations of RT-DNA.
69. Das K, Sarafianos SG, Clark AD Jr, Boyer PL, Hughes SH, Arnold E: **Crystal structures of clinically relevant Lys103Asn/Tyr181Cys double mutant HIV-1 reverse transcriptase in complexes with ATP and non-nucleoside inhibitor HBY 097.** *J Mol Biol* 2007, **365**:77-89.

70. Kleim JP, Bender R, Kirsch R, Meichsner C, Paessens A, Rosner M, Rubsamens-Waigmann H, Kaiser R, Wichers M, Schneeweis KE *et al.*: **Preclinical evaluation of HBY 097, a new nonnucleoside reverse transcriptase inhibitor of human immunodeficiency virus type 1 replication.** *Antimicrob Agents Chemother* 1995, **39**:2253-2257.
  71. Huang W, Gamarnik A, Limoli K, Petropoulos CJ, Whitcomb JM: **Amino acid substitutions at position 190 of human immunodeficiency virus type 1 reverse transcriptase increase susceptibility to delavirdine and impair virus replication.** *J Virol* 2003, **77**:1512-1523.
  72. Xu HT, Asachop EL, Oliveira M, Quashie PK, Quan Y, Brenner BG, Wainberg MA: **Compensation by the E138K mutation in HIV-1 reverse transcriptase for deficits in viral replication capacity and enzyme processivity associated with the M184I/V mutations.** *J Virol* 2011, **85**:11300-11308.
- The study used enzymatic and cell-based assays to show that E138K mutation restored the viral replication capacity in the presence of M184I/V mutation, and E138K developed moderate resistance to efavirenz.
73. Cohen CJ, Molina JM, Cahn P, Clotet B, Fourie J, Grinsztejn B, Wu H, Johnson MA, Saag M, Supparatpinyo K *et al.*: **Efficacy and safety of rilpivirine (TMC278) versus efavirenz at 48 weeks in treatment-naïve HIV-1-infected patients: pooled results from the phase 3 double-blind randomized ECHO and THRIVE Trials.** *J Acquir Immune Defic Syndr* 2012, **60**:33-42.
- This paper summarizes results from two clinical trials using doses of 25 mg rilpivirine versus 600 mg efavirenz, both in combination with TDF/3TC (also AZT/3TC or ABC/3TC in the THRIVE trial) – 25 mg/day rilpivirine had comparable response to 600 mg/day efavirenz. Rilpivirine was better tolerated, yet had higher virological failure rate than efavirenz.
74. Hsiou Y, Ding J, Das K, Clark AD Jr, Hughes SH, Arnold E: **Structure of unliganded HIV-1 reverse transcriptase at 2.7 Å resolution: implications of conformational changes for polymerization and inhibition mechanisms.** *Structure* 1996, **4**:853-860.
  75. Das K, Bauman JD, Clark AD Jr, Frenkel YV, Lewi PJ, Shatkin AJ, Hughes SH, Arnold E: **High-resolution structures of HIV-1 reverse transcriptase/TMC278 complexes: strategic flexibility explains potency against resistance mutations.** *Proc Natl Acad Sci U S A* 2008, **105**:1466-1471.
  76. Udier-Blagovic M, Tirado-Rives J, Jorgensen WL: **Structural and energetic analyses of the effects of the K103N mutation of HIV-1 reverse transcriptase on efavirenz analogues.** *J Med Chem* 2004, **47**:2389-2392.
  77. Ren J, Nichols CE, Stamp A, Chamberlain PP, Ferris R, Weaver KL, Short SA, Stammers DK: **Structural insights into mechanisms of non-nucleoside drug resistance for HIV-1 reverse transcriptases mutated at codons 101 or 138.** *FEBS J* 2006, **273**:3850-3860.
  78. Das K, Bauman JD, Rim AS, Dharia C, Clark AD Jr, Camarasa MJ, Balzarini J, Arnold E: **Crystal structure of tert-butylidimethylsilyl-spiroaminooxathiole-dioxide-thymine (TSAO-T) in complex with HIV-1 reverse transcriptase (RT) redefines the elastic limits of the non-nucleoside inhibitor-binding pocket.** *J Med Chem* 2011, **54**:2727-2737.
  79. Singh K, Marchand B, Rai DK, Sharma B, Michailidis E, Ryan EM, Matzek KB, Leslie MD, Hagedorn AN, Li Z *et al.*: **Biochemical mechanism of HIV-1 resistance to rilpivirine.** *J Biol Chem* 2012, **287**:38110-38123.
- The study used two transient kinetic (quench-flow and stopped-flow) techniques to analyze the impacts of E138K and M184I mutations on dNTP binding, nucleotide incorporation, and the binding of rilpivirine, an NNRTI drug. The analysis revealed that E138K reduces binding affinity of rilpivirine by altering the rate of dissociation versus association of rilpivirine with RT.
80. Selmi B, Deval J, Alvarez K, Boretto J, Sarfati S, Guerreiro C, Canard B: **The Y181C substitution in 3'-azido-3'-deoxythymidine-resistant human immunodeficiency virus, type 1, reverse transcriptase suppresses the ATP-mediated repair of the 3'-azido-3'-deoxythymidine 5'-monophosphate-terminated primer.** *J Biol Chem* 2003, **278**:40464-40472.
  81. Li JZ, Paredes R, Ribaudo HJ, Svarovskaia ES, Metzner KJ, Kozal MJ, Hullsiek KH, Balduin M, Jakobsen MR, Geretti AM *et al.*: **Low-frequency HIV-1 drug resistance mutations and risk of NNRTI-based antiretroviral treatment failure: a systematic review and pooled analysis.** *JAMA* 2011, **305**:1327-1335.
- An analysis of data from 10 clinical studies indicated that the patients carrying low-frequency NNRTI-resistance mutations have an increased risk of virologic failure with first-line antiretroviral treatment (ART).
82. Malcolm RK, Fetherston SM, McCoy CF, Boyd P, Major I: **Vaginal rings for delivery of HIV microbicides.** *Int J Womens Health* 2012, **4**:595-605.
  83. Johnson VA, Calvez V, Gunthard HF, Paredes R, Pillay D, Shafer R, Wensing AM, Richman DD: **2011 update of the drug resistance mutations in HIV-1.** *Top Antivir Med* 2011, **19**:156-164.
  84. Molina JM, Cahn P, Grinsztejn B, Lazzarin A, Mills A, Saag M, Supparatpinyo K, Walmsley S, Crauwels H, Rimskey LT *et al.*: **Rilpivirine versus efavirenz with tenofovir and emtricitabine in treatment-naïve adults infected with HIV-1 (ECHO): a phase 3 randomised double-blind active-controlled trial.** *Lancet* 2011, **378**:238-246.